Project	Description	Elements	Measurement	Length	Status/Results
1 Canadian Nurses Association's Guide to Preceptorship and Mentoring Report Online: http://www.cna.aiic.ca/CNA/nursing/education/mentorship/default_e.aspx Entitled, "Achieving Excellence in Professional Practice" **International**	General guide for setting up a mentoring and preceptorship for novice nurses.	Relevant terms defined Benefits cited Costs explored Steps for developing a successful program identified Preceptor/mentoring competencies identified	Reviewed literature		

Printed from www.ncsbn.org – National Council of State Boards of Nursing, Inc

Project	Description	Elements	Measurement	Length	Status/Results
7 AHRQ: Medical Errors: The Scope of the Problem: An Epidemic of Errors Report from Agency for Healthcare Research and Quality, retrieved February 23, 2009, from: http://www.ahrq.gov/qual/errback.htm **National**	Publication No. AHRQ 00-P037	N/A	Summary of reports on national governmental data.	N/A	Errors occur in settings other than hospitals, including physician's offices, nursing homes, pharmacies, urgent care centers, and care delivered at home. For example, investigations from the MA State Board of Registration in Pharmacy estimate that 2.4 million prescriptions are filled improperly each year in that state. Medical errors cost the nation approximately $36 billion annually, with about $17 billion being related to preventable errors. From IOM 1999 report "To Err is Human: Building A Safer Health System," 44,000 to 98,000 people die each year from medical errors. According to a national poll, 42% of respondents have been affected by a medical error, either personally or through a friend or relative; 32% of the respondents indicate the error had a permanent negative effect on the patient's health; respondents rated the health care system as moderately safe (4.9 on a scale of 1-7), with 7 being "very safe." In another survey, Americans are "very concerned" about being given the wrong medication (61%); being given medications that negatively interact (58%). A landmark study on medical errors found that 70% were preventable; another study showed that 54% of surgical errors were preventable.

Project	Description	Elements	Measurement	Length	Status/Results
10 Carnegie study Some information about the study on their Web site: http://www.carnegiefoundation.org/ A partial report was published here: Benner, P., Sutphen, M. & Leonard-Kahn, V. (2008). Formation and everyday comportment. *A J N C C*, 17(5), 473-476. **National**	Part of larger, national study. Research design was qualitative ethnography, utilizing interviews (total of 588 individual interviews), focus groups, review of curricula, and observations in the classroom and clinical facilities, in excellent nursing programs. Furthermore 3 national surveys were conducted with members of the American Association of Colleges of Nursing, the National League of Nursing, and the National Student Nurse Association.	Recommendation 9.b. states: We recommend residency training programs lasting at least one year focused on one area of nursing care to be offered in all health care delivery institutions. Residency should focus on at least one area of specialty so the nurse has the opportunity to develop in-depth clinical patient population knowledge in that area. Improved follow-through			

Printed from www.ncsbn.org – National Council of State Boards of Nursing, Inc

Printed from www.ncsbn.org – National Council of State Boards of Nursing, Inc

Project	Description	Elements	Measurement	Length	Status/Results
13 Hofler, L. D. 2008. Nursing education and transition to the work environment: A synthesis of national reports. *J [illegible] N[illegible] E [illegible]*, 47(1), 5-12. **National**	2-part process to identify reports and to analyze their content. First organizations were identified (using experts), and then each site was used to retrieve and analyze their work. They purposely did not include regulatory agencies and NCSBN because "their mission is to protect the public." They identified 15 organizations and 35 reports.	Reports identified were between 1995-2005. For inclusion, each report: Was published by a nursing professional organization. Included recommendations			

Project	Description	Elements	Measurement	Length	Status/Results
18 NCSBN Employer Survey (2004) Research Brief is available: https://www.ncsbn.org/360.htm Vol. 14, "Report of Findings from the 2003 Employers Survey" **National**	Surveys completed by 1,230 employers from all settings.	N/A	Survey was investigator constructed.	N/A	Employers answered "Yes definitely" to overall preparation to provide safe, effective care: ADN – 41.9% (n=321) BSN – 41.9% (n=239) Diploma – 48.8% (n=106) LPN – 32.9% (n=237)
19 NCSBN Interim Results of Post-Entry Study: Preliminary report available from NCSBN. Full report is due in September 2008. **National**	Longitudinal, qualitative study of new nurses with 1,111 e-mail responses to date. LPN responses not coded yet.	N/A	Email responses with qualitative analysis about how competence develops	N/A	Implications for transition to practice: The diversity of practice settings and extreme acuity of hospital settings suggest a site-specific transition program with a preceptor for the first year. The narratives demonstrated a real need for novice nurses to revisit action and decisions and reflect on alternate pathways (i.e., need to debrief and reflect). Need for role clarification relative to LPNs and PCAs. Supervision of LPNs or PCAs was either minimal or totally absent.
20 NCSBN's Transition Study (2006) Research Brief available at: https://www.ncsbn.org/360.htm Vol. 22 – "Report of Findings from the Practice and Professional Issues Survey; Transition to Practice: Newly Licensed RN and LPN/VN Activities." April 2006 **National**	NCSBN conducted a survey on 628 new nurses and 519 new LPNs related to transition to practice issues. Survey was investigator constructed.	N/A	Survey was investigator constructed.	N/A	LPNs assigned to care for patients earlier and caseload heavier 38.9% of RNs participated in "ships" + orientation 16.2% of LPNs participated in "ships" + orientation Graduates of ADN programs were more likely than BSN graduates not to have a "ship" Across nation, transition programs were quite variable

Project	Description	Elements			

Project	Description	Elements	Measurement	Length	Status/Results
22 NCSBN data presented at a national forum, entitled "Transition of New Nurses to Practice: A Regulatory Perspective," in Chicago, February 22, 2007. **National**- RNs	*f*N=560 new nurses; N=231 preceptors Non-experimental, compara				

Project	Description	Elements	Measurement	Length	Status/Results
24 Smith, J. & Crawford, L. (2003). Medication errors and difficulty in first patient assignments of newly licensed nurses. *JONA' H L , E ,* , 5(3), 65-67. **National**	This was an NCSBN national study new graduate RNs and LPNs with a focus on medical errors. The surveys were sent to stratified random samples of 1000 RNs (65.5% return rate) and 1000 LPNs/VNs (62.3% return rate).	N/A	Investigator designed tool with new nurse self reports.	N/A	40% of the new LPN/VN graduates were employed in long-term care facilities, with 38% in hospitals and 17% in community or ambulatory care settings. 87% of the new RN graduates were employed in hospitals, with 6% in long-term care facilities and 4% in community or ambulatory care settings. 63% of the new RN graduates were employed in urban/ metropolitan areas, while 47% of the new LPN/VN graduates were employed in urban/ metropolitan areas. 49% of the new RN graduates and 41% of the new LPN/VN graduates made errors or were involved in errors. Of the errors, 75% of the new RN graduates and 71% of the new LPNs/VN graduates were involved with medication errors. Forty percent of the new RN graduates and 47% of the new LPN/VN graduates were involved with errors related to patient falls. Some of the reasons for errors included inadequate staffing (74% of the new LPN/VN graduates and 70% of the new RN graduates), communication (44% of new RN graduates and new LPN graduates), and inadequate orientation (27% of the new LPN/VN graduates and 18% of the new RN graduates. In hospitals new RN graduates cared for an average of 3 patients in their first assignment, and that occurred on an average of 8 days after being hired. New LPN graduates cared for an average of 4 patients and that occurred on an average of 6 days after being hired. In nursing homes new RN graduates averaged 25 patients at the start, whereas LPNs/ VNs cared for an average of 26 patients on their first assignment.

Project	Description	Elements	Measurement	Length	Status/Results
25 University HealthSystem Consortium/American Association of Colleges of Nursing (UHC/AACN): Williams, C.A., Goode, C.J., Krsek, C., Bednash, G.D. and Lynn, M. R. (2007). Postbaccalaureate nurse residency 1-year outcomes. *JONA*. 37(7/8), 357-365. **National**	This is another national, standardized model that is being implemented in 34 sites in university healthcare settings in 24 states.	Core curriculum with focus on leadership, research based practice, professional development, communication, critical thinking, patient			

Project	Description	Elements	Measurement	Length	Status/Results
27 California Institute for Nursing and Health Care Information available here: http://www.cinhc.org/ **Statewide**	A collaborative project in California where they are working to redesign nursing education. Their work groups include: Academic/Service Partnerships; Professional and Clinical Role Development;				

Project	Description	Elements	Measurement	Length	Status/Results
29 Massachusetts Department of Public Health Board of Registration in Nursing: "A Study to Identify Evidence-Based Strategies for the Prevention of Nursing Errors" – Preliminary Data Report available from NCSBN. **Statewide**	Descriptive study of nursing errors found in 78 complaint cases involving 34 RNs and 44 LPNs who practiced in nursing homes in Massachusetts; sampling technique was presented.	N/A	Used a case analysis format, with data being collected using a modified Taxonomy of Error, Root Cause Analysis and Practice Responsibility (TERCAP™) audit instrument.		Seven of the 44 LPNs were licensed for 12 months or less; there were no novice RNs in the analysis. Applicable to the Transition to Practice initiative, errors were linked to inexperience to particular clinical events; lack of familiarity with the practice setting; lack of consistently assigned preceptors and the adequacy of the novice nurse's transition program. Interruptions challenged the novice LPNs who made errors, thus effecting their organizational, prioritizing, communication, delegation, and task completion skills. Study calls attention to the potential patient safety benefit of a novice nurse transition program that provides sufficient time, supervision, and support to new nurses.
30 Mississippi Office of Nursing Workforce Nurse Residency Program Information available: www.monw.org **Statewide**	6-month residency/ internship program, which is implemented through the Mississippi Office of Nursing Workforce.	Coordinator Weekly meetings/			

Project	Description	Elements	Measurement	Length	Status/Results
31 North Carolina's Transition program Information is available here: http://www.ffne.org/transition.cfm **Statewide**	Long-range goal is to create a regulatory model for transitioning new nurses in NC by 2015. Our Research Department is using our transition tools, so these results should enrich our 2006 study results.	Phase I – studying the current transition practices and their impact on newly licensed nurses. Phase II – will focus on developing evidence-based, population-specific transition programs for NC.	Phase I NCSBN's Clinical		

Project	Description	Elements	Measurement	Length	Status/Results
33 Wisconsin Nurse Residency Program (WNRP) Information available here: http://wnrp.org/ **Statewide**	Statewide with 40 plus hospitals, including a large rural group, which is a unique aspect of this program. They have enrolled over 300 new graduates in this program.	Clinical coach Learn to think like a professional Meet once a month Reflection and feedback Focus on: Critical thinking Systems Failure to rescue Best practice EBP Delegation Communication	They look at job stress, organization commitment, clinical decision-making, and behavior in the professional role. Tools include: Porter and Steers Organizational Commitment Jenkins's clinical decision-making Professional Nursing Behavior	12 months	Just finished 3 year HRSA report and have a grant for another 3 years. Are looking to possibly collaborate with NCSBN on use of our transition tool. Will focus on preceptors this time. Increase of retention; rural settings found it highly beneficial.
34 Bjørk, I.T. and Kirkevold, M. (1999). Issues in nurses' practical skill development in the clinical setting. Journal of Nursing Care Quality. 14(1), 72-84. **Individual**	Longitudinal, videotaped interviews of 4 nurses from 8-14 months after licensure Interviews with patients and nurses Practicing skills of dressing changes and ambulation	N/A	Videotapes Interviews with patients and nurses	Had short orientation of 3 weeks	While the nurses became more efficient, they made the same omissions after 14 months: Contaminated wounds Misuse of gloves Failed to wash hands Dangerous tube removal Interviews with patients showed caring over the year Inadequate physical support during ambulation Privacy not maintained Conclusion: Limited orientation/transition program did not allow for reflection and/or feedback so that the same errors were made. Results are relevant for regulation and public protection.

Project					

Project	Description	Elements	Measurement	Length	Status/Results
39 Elfering, A., Semmer, N.K. & Grebner, S. (2006). Work stress and patient safety: Observer-rated work stressors as predictors of characteristics of safety-related events reported by young nurses. *E* , 49(5-6). 457-469. **Individual**	Study conducted in 19 hospitals in Switzerland for a period of 2 working weeks on 23 novice nurses (first 18 months). Stressful events were recorded by the novice nurses, and chronic work characteristics were rated by trained observers, thus providing more validity to this study than those with only self-reports.	N/A	Participants were instructed to document every stressful situation they experienced, requiring an open-ended discussion followed by quantitative items. Compliance with safety regulations was measured with one item, and observers rated chronic job stressors and control on the Instrument for Stress Oriented Task Analysis. Observations were complemented by interviews with the employee, supervisors, and colleagues and consulting with organization documents, if necessary.	N/A	62 events, or 2.65 events per person, were related to patient safety Safety events included: documentation, near misses with medication, incomplete patient briefing, delays in care, patient casualties (falls, etc.). Stressors, most notably concentration demands and lack of control, related endangered patient safety. Recommendation: training of novice nurses should address the association between workload and patient safety and should education nurses in self-management strategies for stress.

Project	Description	Elements	Measurement	Length	Status/Results
41 Johnstone, M. J. and Kanitsaki, O. (2006). Processes influencing the development of graduate nurse capabilities in clinical risk management: An Australian study. **Individual**	Exploratory-descriptive case study approach, with qualitative and quantitative data collection and analysis	Elements (from literature) of opportunities putting graduate nurses at risk for error: Inadequate education Inadequate supervision Workplace bullying Hierarchical structures inhibiting performance Poor planning and scheduling of work Poor skill mix Heavy workload			

data:/MCID 3 /Artifact /Lang (en) /BBox [70 818 24 L /3R3 gs/C2_0 1 Tf8 0 0 8 285.7651 464.519 Tm[0033]TjETEMC /Sd analysisD 323litative and quantitative
Inquiry 2014 descriptive

Project	Description	Elements	Measurement	Length	Status/Results
44 Latham, C., Hogan, M., Ringl, K. (2008). Nurses supporting nurses: Creating a mentoring program for staff nurses to improve the workforce environment. *N[illegible] A[illegible]*, 32(1), 27-39. **Individual**	A 3-year academic-hospital partnership was developed to create a mentoring program for nurses to improve the workplace environment: Nurses Supporting Nurses. This program was targeted for new graduates and new hires, though any staff RN was welcome to apply for the mentee role. Ninety two mentor-mentee teams were formed. The program was particularly focused on changing the culture of the workplace. Two hospitals with quite different organizational characteristics were utilized for the project. The purpose was to identify future bedside leaders who would assume supportive roles, thereby changing the culture of the unit.	Hospital liaison to champion the project, and this liaison was key to the success. Interactive workshops on culture mindedness, which included team building. Creation of detailed Web pages, including videoclips of mentors. Sociometric analysis of RN camaraderie and informal leadership. Educational sessions on nurses supporting nurses. Cultural competence was presented. Speed meetings to help select mentors. Other sessions included team building, conflict resolution, communication, time management, healthcare system, financial concerns, quality care, patient satisfaction, and safety. Quarterly mentor support meetings.	Overall nurse satisfaction Decisional involvement Cultural communication competency Retention and vacancy Patient satisfaction with nursing care 3 nurses sensitive areas, including falls, pressure ulcer prevention, and proper use of restraints	3-year	Personality and learning styles are not the basis of successful mentor-mentee teams. Most nurses believed they were culturally competent and that the environment supported cultural sensitivity. Most nurses wanted more control over their working conditions. Improvements in patient and nurse satisfaction. Improvement in nurse vacancy and turnover. Improvement related to fall and pressure ulcer prevention and on the proper use of restraints. Using $100,000 per RN replacement charge, the 2 hospitals had a cost savings of $2.5 million.

Project	Description	Elements	Measurement	Length	Status/Results
45 Launch into Nursing: a collaboration between the University of Texas M.D. Anderson Cancer Center and The University of Texas Health Science Center at Houston, School of Nursing. Keller, J. L., Meekins, K. and Summers, B.L. (2006). Pearls and pitfalls of a new graduate academic residency program. *JONA.* 36(12), 589-598. **Individual**	Describe the design of a collaborative academic residency program for graduate nurses.	Academic leadership course has become cornerstone. Also included simulations, including "Friday Night in the ER." Introduction to workplace resources, which included projects, small group discussions, introductions to a variety of roles, etc. Socialization was very important, as it has been cited as linked to retention and safety. Each was matched to a trained "clinical coach." Describes their curriculum map in detail. Will be helpful with module design. Areas from curriculum map include: Communication Systems thinking Safety EBP Socialization QI	Outcomes measured, with various tools: Progress to competent nurse (Benner) Knowledge Retention Intent to leave Job satisfaction Employee engagement Competence in clinical leadership Comprehension of Magnet essentials Evidence-based practice techniques Commitment to lifelong learning Culture of support Cultural competency Role as patient advocate Successful acculturation Accountability	12 months	Education, at its best, cannot prepare for acculturation into a work group, using a newly learned language in practice, becoming proficient in a wide range of absolutely necessary skills, and gaining a sense of the wider world of health care. Incorporates reflection and feedback Turnover at 1 year was 10.8% Cost was $1,000 per resident Estimated that cost of replacing 1 nurse was $60,000
46 Merry, M.D. & Brown, J.P. (2001). From a culture of safety to a culture of excellence: Quality science, human factors, and the future of healthcare quality. Journal of Innovative Management , 7(2), 29-46. **Individual**	Report of the sigma gap in health care, which is the gap between performance and potential performance.	N/A	N/A	N/A	Airlines – 0.43 deaths per million passengers Hospitalizations – 2,300 deaths due to error per million admits

Printed from www.ncsbn.org – National Council of State Boards of Nursing, Inc